

Table 13. Estimated proportion of persons surviving for more than 12, 24, and 36 months after an AIDS diagnosis in 1998, by selected characteristics—United States

No. of persons	Survival in months		
	>12	>24	>36
	Proportion		
Age at diagnosis (yrs)			
<13	228	0.91	0.88
13–14	54	0.96	0.94
15–24	1,553	0.95	0.93
25–34	12,160	0.93	0.89
35–44	16,868	0.90	0.86
45–54	7,543	0.86	0.81
55–64	2,039	0.79	0.74
≥65	663	0.72	0.65
Race/ethnicity			
White, not Hispanic	12,998	0.91	0.87
Black, not Hispanic	19,669	0.89	0.84
Hispanic	7,925	0.89	0.86
Asian/Pacific Islander	330	0.90	0.89
American Indian/Alaska Native	151	0.90	0.85
Exposure category			
Male adult or adolescent			
Male-to-male sexual contact	14,643	0.91	0.88
Injection drug use	6,645	0.87	0.82
Male-to-male sexual contact and injection drug use	2,122	0.92	0.88
Heterosexual contact	2,945	0.91	0.86
Other ^a	4,728	0.85	0.81
Subtotal	31,083	0.90	0.86
Female adult or adolescent			
Injection drug use	2,945	0.89	0.82
Heterosexual contact	4,563	0.92	0.87
Other ^a	2,287	0.88	0.85
Subtotal	9,795	0.90	0.85
Child(<13 yrs)			
Perinatal	221	0.91	0.88
Other ^b	7	0.86	0.86
Subtotal	228	0.91	0.88
Total^c	41,108	0.90	0.86
0.82			

Note. Excludes persons whose date of death is before, or in the same month as, date of diagnosis. Excludes persons whose month of diagnosis or month of death is unknown.

^a Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk not reported or not identified.

^c Includes 35 persons of unknown or multiple race and 2 persons of unknown sex.